



Complete Circular Genome Sequences of *Brachyspira hyodysenteriae* Isolates of the Four Different Sequence Types Causing Swine Dysentery in Switzerland

 Ana B. García-Martín,^{a,b} Sarah Schmitt,^c Friederike Zeeh,^d  Vincent Perreten^a

^aDivision of Molecular Epidemiology and Infectious Diseases, Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Bern, Switzerland

^bGraduate School for Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland

^cInstitute for Food Safety and Hygiene, Section of Veterinary Bacteriology, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland

^dClinic for Swine, Department of Clinical Veterinary Medicine, Vetsuisse Faculty, University of Bern, Bern, Switzerland

ABSTRACT The complete genomes of four *Brachyspira hyodysenteriae* isolates of the four different sequence types (STs) (ST6, ST66, ST196, and ST197) causing swine dysentery in Switzerland were generated by whole-genome sequencing and *de novo* hybrid assembly of reads obtained from second (Illumina) and third (Oxford Nanopore Technologies and Pacific Biosciences) generation high-throughput sequencing.

B *Brachyspira hyodysenteriae* was confirmed to cause swine dysentery in Swiss pig herds in 2008 (1). Since then, isolates have been characterized by multilocus sequence typing and determination of their antimicrobial resistance mechanisms (2, 3). To date, only four sequence types (STs) (ST6, ST66, ST196, and ST197) of *B. hyodysenteriae* have been identified, prompting us to sequence their complete genomes.

Isolates obtained from our cryopreserved collections in Zurich (BHZ755 [ST6], BHZ375 [ST66], and BHZ526 [ST197]) and Bern (Bh743-7 [ST196]) were grown anaerobically at 42°C for 5 days on Trypticase soy agar with 5% (vol/vol) defibrinated sheep blood (Becton, Dickinson). The bacterial lawn of two plates was collected using a 10- μ l plastic loop and resuspended in buffer (0.1 M Tris-HCl, 0.01 M NaCl, 0.1 M EDTA). Genomic DNA was extracted using the DNeasy blood and tissue kit (Qiagen), treated with RNase (20 mg/ml) at 37°C for 30 min, purified with AMPure XP magnetic beads (Beckman Coulter), and quantified with a Qubit 3.0 fluorometer (Life Technologies). Prior to Oxford Nanopore Technologies (ONT) long-read library preparation, DNA was sheared up to 20 kb using Covaris g-TUBEs. DNA libraries were obtained using the 1D native barcoding (EXP-NBD104) and ligation (SQK-LSK109) kits and loaded onto a FLO-MIN106D flow cell R9.4.1. Real-time sequencing was performed using a MinION Mk1B device (ONT), and results were visualized using the controlling software MINKNOW-GUI v19.05.0 (ONT). Base calling and demultiplexing were performed using Guppy v2.3.7 (ONT). Long reads were analyzed using NanoPack (4) and trimmed by 100 bp using Cutadapt v1.18 (5). Default parameters were used for all software unless otherwise specified.

Pacific Biosciences (PacBio) long-read libraries were prepared from DNA extracted as described previously (6), following the BluePippin \geq 7-kb size selection method (with an insert length of 10 kb), and sequenced in a single-molecule real-time (SMRT) Cell v2 compatible with the Sequel II system at Lausanne Genomic Technologies Facility (Lausanne, Switzerland). Reads were demultiplexed using lima from the SMRT Analysis software (PacBioSuite-6.0.0.47841) (<https://www.pacb.com/support/software-downloads>) and analyzed using LongQC (7).

Short-read sequencing was performed using a NEBNext Ultra II directional DNA library with TruSeq adapters on an Illumina NovaSeq 6000 platform at Eurofins (Germany). Paired-end 2 \times 150-bp reads were quality checked using FastQC v0.11.7 (8) and filtered

Citation García-Martín AB, Schmitt S, Zeeh F, Perreten V. 2021. Complete circular genome sequences of *Brachyspira hyodysenteriae* isolates of the four different sequence types causing swine dysentery in Switzerland. Microbiol Resour Announc 10:e00847-21. <https://doi.org/10.1128/MRA.00847-21>.

Editor Steven R. Gill, University of Rochester School of Medicine and Dentistry

Copyright © 2021 García-Martín et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Vincent Perreten, vincent.perreten@vetsuisse.unibe.ch.

Received 24 August 2021

Accepted 28 August 2021

Published 30 September 2021

TABLE 1 Metrics of the announced complete genomes of *Brachyspira hyodysenteriae* isolates

Parameter	Data for isolate:			
	Bh743-7	BHZ375	BHZ526	BHZ755
Isolate information				
Isolation date	2017	2013	2014	2015
ST	196	66	197	6
Raw sequencing data				
ONT^a				
No. of sequenced reads (Unicycler-loaded reads)	346,463	220,134	313,725	276,045
Sequenced length (bp)	1,744,406,639	819,961,706	1,859,824,743	1,504,066,585
Mean read length (bp)	5,235	3,925	6,128	5,649
Read N_{50} (bp)	7,705	6,528	8,858	8,264
PacBio^b				
No. of sequenced reads	189,845	177,937	170,348	186,892
No. of Unicycler-loaded reads	189,444	177,933	170,328	186,839
Sequenced length (bp)	NA	NA	NA	NA
Mean read length (bp)	7,037	7,229	6,940	6,984
Read N_{50} (bp)	9,048	9,163	9,056	8,898
Illumina^c				
No. of sequenced reads ($\times 10^6$)	7.4	10.8	6.3	6.2
Sequenced length (Mbp)	2,225	3,243	1,879	1,870
Assembly^d				
ONT-Illumina				
Total genome length (bp)	3,085,162	3,043,241	3,036,451	3,035,657
Chromosome length (bp)	3,052,634	3,010,106	3,003,989	3,003,105
Plasmid length (bp)	32,528	33,135	32,462	32,552
G+C content (%)	27.0	27.0	27.1	27.1
Coverage depth (mean \pm SD) (\times)				
Chromosome	691 \pm 84	1,001 \pm 119	590 \pm 51	592 \pm 56
Plasmid	1,203 \pm 118	3,510 \pm 319	1,004 \pm 97	1,036 \pm 101
PacBio-Illumina				
Total genome length (bp)	3,085,111	3,043,198	3,035,137	3,035,627
Chromosome length (bp)	3,052,583	3,010,063	3,002,675	3,003,075
Plasmid length (bp)	32,528	33,135	32,462	32,552
G+C content (%)	27.1	27.0	27.1	27.1
Coverage depth (mean \pm SD) (\times)				
Chromosome	691 \pm 84	1,001 \pm 119	590 \pm 53	592 \pm 56
Plasmid	1,203 \pm 118	3,510 \pm 319	1,004 \pm 97	1,036 \pm 101
ONT-Illumina hybrid assembly correction^e				
No. of Sanger sequencing-corrected regions	3	1	1	2
Total no. of nucleotides added per chromosome	59	15	9	27
Primer sequences (5' to 3')				
Chromosomal region 1 (16 bp) (nucleotide positions 444843 to 444858)	Bh743-7-1F, AGTACCTTTCCAGCAGCAAG; Bh743-7-1R, GCAGAGGTGAAGCCGCTAAA			
Chromosomal region 2 (7 bp) (nucleotide positions 449527 to 449533)	Bh743-7-2F, TGTACAAAATTATATGCCGCCATAA; Bh743-7-2R, TTGATGAGCGTATGTGGGAAT			
Chromosomal region 3 (36 bp) (nucleotide positions 2726169 to 2726204)	Bh743-7-3F, GGAAATAGTTGGGGAGAGTTCA; Bh743-7-3R, TGGAAATAGTAAACCCAAATCTGTTG			
Chromosomal region 1 (15 bp) (nucleotide positions 1728861 to 1728875)	BHZ375-2F, CCCCATAAAAAGCTTTGAATCCA; BHZ375-2R, TGCTATGCAGATGCGTTTGC			
Chromosomal region 1 (9 bp) (nucleotide positions 2678040 to 2678048)	BHZ526-1F, AGGAAATAGTTGGGGAGAGGT; BHZ526-1R, AGCAAGATGATGTTATGCTGT			
Chromosomal region 1 (20 bp) (nucleotide positions 448627 to 448782)	Bh743-7-1F, AGTACCTTTCCAGCAGCAAG; Bh743-7-1R, GCAGAGGTGAAGCCGCTAAA			
Chromosomal region 2 (7 bp) (nucleotide positions 1283746 to 1283752)	BHZ755-2F, TGAGCATAAGCACGGCATT; BHZ755-2R, ACTAATGCTGTATCTCCAATCCA			
Final ONT-Illumina hybrid assembly annotation^f				
Total no. of coding sequences	2,652	2,590	2,588	2,595
No. of coding genes	2,623	2,557	2,558	2,566
Total no. of rRNAs	40	40	40	40
No. of pseudogenes	29	33	30	29

(Continued on next page)

TABLE 1 (Continued)

Parameter	Data for isolate:			
	Bh743-7	BHZ375	BHZ526	BHZ755
Accession numbers ^a				
SRA database				
ONT	SRR10609650	SRR10609648	SRR10609646	SRR10609644
Illumina	SRR10609649	SRR10609647	SRR10609645	SRR10609643
PacBio	SRR15444232	SRR15444231	SRR15444230	SRR15444229
GenBank nucleotide database				
Chromosome (ONT-Illumina hybrid assembly)	CP046932	CP046930	CP046928	CP046926
Plasmid (ONT-Illumina hybrid assembly)	CP046933	CP046931	CP046929	CP046927

^aObtained with NanoPack (Nanostat) and LongQC.

^bObtained with LongQC. NA, not applicable.

^cObtained with FastQC.

^dObtained with QUAST and Illumina short-read remapping by running BMap in Geneious.

^eNumbers of regions and nucleotides that were manually inserted per chromosome in each genome after PCR and Sanger sequencing confirmation. Chromosomal region 1 of the BHZ755 genome was corrected using primers Bh743-7-1F and Bh743-7-1R.

^fObtained from the NCBI PGAP annotation files.

^gAccession numbers corresponding to the sequencing runs deposited in the SRA database and accession numbers of the nucleotide sequences deposited in the GenBank sequence database.

with Trimmomatic v0.36 (9). ONT and Illumina libraries were prepared from the same DNA samples, and PacBio libraries were prepared from different batches of DNA obtained using the same culture conditions.

De novo hybrid assemblies were generated by running the *-bold* option of Unicycler v0.4.4 (10). Assembly polishing, circularization, and rotation were performed using Unicycler. Polished circular assemblies were visualized using Bandage v0.8.1 (11). Assembly quality was analyzed using QUAST v4.6.0 (12). The mean depth of coverage (Table 1) for the entire assembly was calculated by remapping Illumina reads by running the plugin BMap v37.25 (13) in Geneious R10.2.3 (Biomatter, Ltd.). Additional comparisons of the hybrid ONT-Illumina assemblies with their PacBio counterparts allowed the detection of chromosomal low-complexity regions, which were corrected by Sanger sequencing (Table 1). Sanger sequencing samples were obtained from the same DNA as the ONT and Illumina libraries and were sequenced in both directions. Sequence chromatograms were visualized in Geneious to assess their quality. Untrimmed sequences were aligned to their corresponding assemblies, which were manually edited in Geneious. Assemblies consisted of two circular replicons (both characterized by low G+C contents of <27.1%), representing one chromosome and one plasmid, with mean lengths of 3,017,459 bp and 32,669 bp, respectively. Annotation was performed using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.10 (14). On average, 2,576 protein-coding genes, 40 rRNAs, and 30 pseudogenes were identified on the chromosome (Table 1). Thirty protein-coding genes were identified in all plasmids except pBHZ375, which has 29 protein-coding genes.

The hybrid approach applied here allowed reconstruction of high-quality genomes (in terms of completeness and accuracy). These genomes contribute to expanding the catalogue of *B. hyodysenteriae* genomes and will serve as a basis for molecular epidemiology studies of swine dysentery.

Data availability. Annotated genomes have been deposited in GenBank under the BioProject accession number [PRJNA594292](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA594292) and BioSample accession numbers [SAMN13511716](https://www.ncbi.nlm.nih.gov/biosample/SAMN13511716), [SAMN13511717](https://www.ncbi.nlm.nih.gov/biosample/SAMN13511717), [SAMN13511718](https://www.ncbi.nlm.nih.gov/biosample/SAMN13511718), and [SAMN13511719](https://www.ncbi.nlm.nih.gov/biosample/SAMN13511719), and their accession numbers are listed in Table 1. ONT and Illumina raw data sets are archived in the SRA database (Table 1). Demultiplexed PacBio raw data (BioProject accession number [PRJNA754405](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA754405)) are also archived in the SRA database (Table 1).

ACKNOWLEDGMENTS

This project was financed by grants 1.16.04 and 1.19.05 from the Swiss Federal Food Safety and Veterinary Office (SFVO) and by internal funds of the Institute of Veterinary Bacteriology of the University of Bern.

We thank the Competency Center in Bioinformatics and Computational Biology (Vital-IT) (<https://www.vital-it.ch>) of the Swiss Institute of Bioinformatics (University of Lausanne, Lausanne, Switzerland) and the high-performance computing cluster (UBELIX) (<https://hpc-unibe-ch.github.io>) of the University of Bern for providing software support and computing resources. We also thank Alexandra Collaud for guidance during ONT library preparation and MinION sequencing.

REFERENCES

- Speiser S. 2008. Untersuchungen zu *Lawsonia intracellularis*, *Brachyspira hyodysenteriae* und *Brachyspira pilosicoli* Infektionen bei Absetz und Mast Schweinen in der Schweiz. Dr. med. vet. dissertation. Vetsuisse Faculty, University of Bern, Bern, Switzerland.
- Kirchgässner C, Schmitt S, Borgström A, Wittenbrink MM. 2016. Antimicrobial susceptibility of *Brachyspira hyodysenteriae* in Switzerland. *Schweiz Arch Tierheilkd* 158:405–410. <https://doi.org/10.17236/sat00066>.
- García-Martín AB, Perreten V, Rossano A, Schmitt S, Nathues H, Zeeh F. 2018. Predominance of a macrolide-lincosamide-resistant *Brachyspira hyodysenteriae* of sequence type 196 in Swiss pig herds. *Vet Microbiol* 226: 97–102. <https://doi.org/10.1016/j.vetmic.2018.10.007>.
- De Coster W, D'Hert S, Schultz DT, Cruys M, Van Broeckhoven C. 2018. NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics* 34:2666–2669. <https://doi.org/10.1093/bioinformatics/bty149>.
- Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 17:10–12. <https://doi.org/10.14806/ej.17.1.200>.
- Tapia-Paniagua ST, Chabrillon M, Díaz-Rosales P, de la Banda IG, Lobo C, Balebona MC, Morinigo MA. 2010. Intestinal microbiota diversity of the flat fish *Solea senegalensis* (Kaup, 1858) following probiotic administration. *Microb Ecol* 60:310–319. <https://doi.org/10.1007/s00248-010-9680-z>.
- Fukasawa Y, Ermini L, Wang H, Carty K, Cheung MS. 2020. LongQC: a quality control tool for third generation sequencing long read data. *G3 (Bethesda)* 10:1193–1196. <https://doi.org/10.1534/g3.119.400864>.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. Babraham Bioinformatics, Cambridge, United Kingdom. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Wick RR, Schultz MB, Zobel J, Holt KE. 2015. Bandage: interactive visualization of *de novo* genome assemblies. *Bioinformatics* 31:3350–3352. <https://doi.org/10.1093/bioinformatics/btv383>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Bushnell B. 2014. BMAP: a fast, accurate, splice-aware aligner. Lawrence Berkeley National Laboratory, Berkeley, CA.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.